

AD-A131 564

TRANSIENT PHOTOCHEMISTRY OF NEUTRAL RED(U) RENSSELAER  
POLYTECHNIC INST TROY N Y DEPT OF CHEMISTRY  
G T MARKS ET AL. JUL 83 TR-5-ONR N00014-78-C-0494

1/1

UNCLASSIFIED


F/G 7/3

NL

END

FORMED

AD



MICROCOPY RESOLUTION TEST CHART  
NATIONAL BUREAU OF STANDARDS-1963-A

12

ADA131564

DTIC FILE COPY

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER ONR-TR-5	2. GOVT ACCESSION NO. AD A131564	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Transient Photochemistry of Neutral Red		5. TYPE OF REPORT & PERIOD COVERED Interim Technical Report
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) G. T. Marks, E. D. Lee, D. A. Aikens and H. H. Richtol		8. CONTRACT OR GRANT NUMBER(s) N0014-78-C-0494
9. PERFORMING ORGANIZATION NAME AND ADDRESS Rensselaer Polytechnic Institute Department of Chemistry Troy, New York 12181		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS NR051-679
11. CONTROLLING OFFICE NAME AND ADDRESS Code 472 Office of Naval Research Arlington, VA 22217		12. REPORT DATE July 1983
		13. NUMBER OF PAGES 34
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report)
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report) B		
18. SUPPLEMENTARY NOTES Prepared for publication in Photochemistry, Photobiology		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) neutral red, transient spectra, semireduced radicals, triplets		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Flash photolysis of neutral red between pH 1.3 and pH 11 yields the triplet species $^3\text{DH}_2^{+2}$ , $^3\text{DH}^+$ , and $^3\text{D}$ . Both $^3\text{DH}_2^{+2}$ , and $^3\text{D}$ exhibit first order decay with rate constants of $1.6 \pm 0.3 \times 10^8 \text{ s}^{-1}$ , but $^3\text{DH}^+$ decays within the lifetime of the flash. Over the entire pH range, ascorbic acid quenches the triplet, forming the semireduced radicals $\text{DH}_3^{+2}$ , $\text{DH}_2^+$ , and $\text{DH}^\cdot$ , all of which exhibit second order decay with $k = 1.8 \pm 0.4 \times 10^8 \text{ M}^{-1} \text{ s}^{-1}$ , most probably by		

recombination with semioxidized ascorbic acid. the dependence of the rate of decay of radical neutral red on the identity of reversible reductants supports the back-electron transfer mechanism, as does digital simulation of complex radical disproportionation schemes. In contrast to the efficient reduction of triplet neutral red by ascorbic acid, its reduction by EDTA is quite inefficient.

OFFICE OF NAVAL RESEARCH  
Contract No. N00014-78-C-0494  
Task No. NR051-679  
Technical Report No. 5

Transient Photochemistry of Neutral Red

by

G. T. Marks, E. D. Lee, D. A. Aikens and H. H. Richtol

Prepared for publication in  
Photochemistry, Photobiology

Rensselaer Polytechnic Institute  
Department of Chemistry  
Troy, New York 12181

July, 1983

Reproduction in whole or in part is permitted for  
any purpose of the United States Government

Approved for Public Release; Distribution Unlimited

# TRANSIENT PHOTOCHEMISTRY OF NEUTRAL RED

by

G. T. Marks, E. D. Lee, D. A. Aikens\*,  
and H. H. Richtol\*

Department of Chemistry  
Rensselaer Polytechnic Institute

Troy, NY 12181

Short Running Title: Transient Photochemistry of Neutral Red

Subheadings  
neutral red

transient spectra

semireduced radicals

triplets

Keywords

transient species, triplets,  
semireduced radicals

neutral red, decay rates, acidity  
constants

formation, decay, influence of  
reducing agents, acidity constants  
acidity constants, decay rates,  
reduction

## ABSTRACT

Flash photolysis of neutral red between pH 1.3 and pH 11 yields the triplet species  $^3\text{DH}_2^{+2}$ ,  $^3\text{DH}^+$ , and  $^3\text{D}$ . Both  $^3\text{DH}_2^{+2}$ , and  $^3\text{D}$  exhibit first order decay with rate constants of  $1.6 \pm 0.3 \times 10^4 \text{ s}^{-1}$ , but  $^3\text{DH}^+$  decays within the lifetime of the flash. Over the entire pH range, ascorbic acid quenches the triplet, forming the semireduced radicals  $\text{DH}_3^{+2}$ ,  $\text{DH}_2^+$  and  $\text{DH}^\cdot$ , all of which exhibit second order decay with  $k = 1.8 \pm 0.4 \times 10^8 \text{ M}^{-1} \text{ s}^{-1}$ , most probably by recombination with semioxidized ascorbic acid. The dependence of the rate of decay of radical neutral red on the identity of reversible reductants supports the back-electron transfer mechanism, as does digital simulation of complex radical disproportionation schemes. In contrast to the efficient reduction of triplet neutral red by ascorbic acid, its reduction by EDTA is quite inefficient.

Accession For	
NTIS	✓
DTIC	
Unannounced	
IN	
DATE	
COPIES	
APPROVED	
Dist	
<b>A</b>	

## INTRODUCTION

Although the transient photochemistry of thiazine dyes such as methylene blue (Wildes et al., 1977) and thionine (Bonneau et al., 1974) has been the subject of intensive study, the transient photochemistry of the azine dyes has received little attention. We report here such a study of neutral red (3-amino-6-dimethylamino-2-methylphenazine hydrochloride). Neutral red has long been used as both an acid-base indicator (Bishop, 1972), and a biological stain (Clark and Perkins, 1932), but its transient photochemistry has been the subject of only one brief study (Chibisov et al., 1969) which reported semireduced radical peaks at pH 4 and pH 11. Because both the triplet species and the semireduced radical species play critical roles in the photoredox reactions of dyes (Ohno and Lichtin, 1980), knowledge of the properties of these excited states is necessary for proper understanding of such processes. The present study establishes the major features of the acid-base behavior, the visible spectra and the decay kinetics of both the semireduced radical species and the triplet species of neutral red. In some respects, the properties of these excited state species of neutral red are similar to those of the thiazine dyes methylene blue (Wildes et al., 1977) and thionine (Bonneau et al., 1974) and the azine dye safranin-O (Baumgartner et al., 1981), but there are important differences which distinguish neutral red from the other dyes.



## MATERIALS AND METHODS

Neutral red was recrystallized thrice from 50% aqueous methanol and other chemicals were reagent grade, and were used without further purification. Except as noted in certain studies with EDTA as reducing agent,  $2 \times 10^{-6}$  M dye in aqueous  $1 \times 10^{-3}$  M phosphate buffer was used throughout. Prior to flash photolysis, the dye solution in a 10 cm quartz or borosilicate cell was rigorously freed of oxygen by 7 successive freeze-pump-thaw cycles under a vacuum of  $10^{-5}$  torr. Spectroscopic transients were secured using the xenon flash photolysis unit described by Strong and Perano (1967), which permitted monitoring the transient absorption between 350 nm and 820 nm beginning 50  $\mu$ s after flash initiation. Before reaching the sample cell, the flash lamp beam passed through a uranyl nitrate solution and a copper sulfate-methylene blue solution to limit the sample illumination to wavelengths between ca 450 nm and 550 nm, the range which corresponds to the absorption band of ground state neutral red. Kinetic data were subjected to linear regression analysis and only data sets with regression coefficients of 0.985 or higher were used to compute rate constants.

## RESULTS AND DISCUSSION

### Triplet State Characteristics

Transient absorption spectra obtained from pH 1.3 to pH 11.0 in the absence of reducing agents, representative examples of which are in Figure 1, were used to estimate the

acidity constants of the triplet state. Strong absorption by ground state dye from ca 440 nm to 550 nm precludes characterization in this range of transient spectra, the essential features of which are as follows. Between pH 1.3 and pH 4.2, the transient absorption spectrum consists of two bands with maxima at 390 nm and 680 nm, the intensities of which are essentially pH independent. Above pH 4.2 the 680 nm band decreases in intensity, the 390 nm band broadens and both peaks shift toward shorter wavelengths. At pH 5.1, the 680 nm peak has shifted to 650 nm and the 390 nm peak has shifted to 380 nm. Accompanying these spectral shifts is a sharp drop in the lifetime of the triplet species, which becomes so short that above pH 6 the triplet absorption follows the decay of the flash and cannot be detected 50  $\mu$ s after flash initiation. Triplet absorption remains undetectable until ca pH 9.2, at which point the triplet lifetime has increased sufficiently to reveal the bands at 380 nm and 650 nm and a new band at 580 nm. Increasing the pH from 9.2 to 11 causes the intensity of the 650 nm band to fall nearly to zero and the intensity of the 580 nm band to increase to a limiting value, while the intensity of the 380 nm band remains essentially constant.

No transients were observed in the presence of oxygen, a result which together with the reactivity of these transient species with reducing agents (see next section) indicates that they are triplet species. Because the  $pK_a$  of neutral red is 6.8 (Bartels, 1956), both the acidic and basic forms of ground

state neutral red give rise to triplet species. From pH 1.3 to pH 4.2 and from pH 9.2 to pH 11.0, the decay rate of the triplet species is fixed, independent of pH. Throughout both pH domains, the transient depletion of the ground state dye shows the triplet concentration 50  $\mu$ s after flash initiation to be  $1.3 \times 10^{-7}$  M. That is, at this point the triplet yield is equivalent to ca 6-7% of the concentration of ground state dye, as compared to ca 50% triplet yield for safranin-O under similar excitation conditions in the same apparatus (Baumgartner et al., 1981). In these studies, the ground state absorption was held approximately constant by centering the irradiation at the ground state isosbestic point of 480 nm. It is noteworthy that over the entire pH range where the triplet can be detected, the triplet yield remains virtually fixed at 6-7%. This estimate of the triplet yield is based on the usual assumption that the triplet species do not absorb significantly at 500 nm, the wavelength at which we monitored the ground state depletion. Although Figure 1 suggests that this assumption is valid, it is possible that the triplet species do absorb at this wavelength. Therefore, the estimates of triplet yield are minimum values, while the estimates of the triplet absorptivities are maximum values.

The molar absorptivity of each triplet species was determined from the ground state depletion and the transient triplet absorbance at a pH which ensures that only one triplet species is present. These values along with the first order

rate constants for triplet decay are reported in Table I. In view of the structural similarity of neutral red to thionine, oxonine and safranin-O, we have adopted the notation of Bonneau *et al.*, (1973) for the triplet state whereby the most highly protonated triplet species in the pH range of interest is denoted  $^3\text{DH}_2^{+2}$ .

The pH dependence of the transient triplet absorption bands indicates that the acidic triplet undergoes stepwise loss of two protons as described by Equation 1.




Figure 2 reports the pH dependence of the normalized absorbance at 680 nm and at 580 nm 50  $\mu\text{s}$  after flash initiation. Although these plots are qualitatively similar to those for a number of other dyes (Knowles, 1971), they cannot be used to determine exact values of the  $\text{pK}_a$ s of  $^3\text{DH}_2^{+2}$  and  $^3\text{DH}^+$ . The reason is that the decay of the intermediate triplet,  $^3\text{DH}^+$ , is much faster than that of  $^3\text{DH}_2^{+2}$ . In fact, it is so rapid that between pH 6 and pH 9 it follows the decay profile of the flash lamp and no triplet absorption can be detected 50  $\mu\text{s}$  after flash initiation. Because  $^3\text{DH}^+$  decays much more rapidly than does  $^3\text{DH}_2^{+2}$ , formation of  $^3\text{DH}^+$  above pH 4 increases the rate of  $^3\text{DH}_2^{+2}$  decay. This coupling occurs because proton transfer is much faster than decay of  $^3\text{DH}_2^{+2}$  (Schulman, 1976), so that throughout its lifetime,  $^3\text{DH}_2^{+2}$

disappears by conversion to  $^3\text{DH}^+$  as well as by direct decay to the ground state. This second decay route lowers the concentration of  $^3\text{DH}_2^{+2}$  throughout its lifetime, and as a result, the pH at which the normalized absorbance of  $^3\text{DH}_2^{+2}$  falls to 0.5 is lower than the  $\text{pK}_a$  of  $^3\text{DH}_2^{+2}$ . The interaction of  $^3\text{DH}^+$  with  $^3\text{D}$  is analogous to its interaction with  $\text{DH}_2^{+2}$ , and similar considerations show that the pH at which the normalized absorbance rises to 0.5 is higher than the  $\text{pK}_a$  of  $^3\text{DH}^+$ . That is, the lower limit of  $\text{pK}_{a_1}$  is 4.8 and the upper limit of  $\text{pK}_{a_2}$  is 9.5. Determination of the precise values of the  $\text{pK}_a$ s requires knowledge of the decay kinetics of  $^3\text{DH}^+$ , which in turn requires that absorption measurements be made sooner after flash initiation. Despite this limitation, however, the estimated  $\text{pK}_a$ s of triplet neutral red (>4.8, <9.5) agree reasonably well with the triplet  $\text{pK}_a$ s of the related dyes thionine (6.3, 8.9) (Fischer, 1964; Faure *et al.*, 1967; Bonneau and Pereyre, 1975) and safranin-O (7.5, 9.2) (Baumgartner *et al.*, 1981).

Both  $^3\text{DH}_2^{+2}$  and  $^3\text{D}$  exhibit strictly first order decay over 2-3 halflives. The absence of a transition to second order kinetics late in the life of the triplet species shows that triplet-ground state interaction forming semireduced and semioxidized dye, which was suspected to occur with safranin-O (Baumgartner *et al.*, 1981), is not significant for neutral red. All transients decay return to the baseline within ca 0.2 ms and the ground state absorption spectrum is unchanged

after samples are subjected to as many as 50 flashes, thus confirming that triplet formation is reversible and that no long-lived species are formed.

#### Semireduced Radical Formation

We have examined the one-electron reduction of triplet state neutral red by several reducing agents, with emphasis on the behavior of ascorbic acid and EDTA. With  $2 \times 10^{-4}$  M ascorbic acid as the reductant, between pH 1.3 and pH 11, the concentration of semireduced radical 50  $\mu$ s after flash initiation estimated from the transient decrease in ground state absorption at 500 nm corresponds to ca 6% of the ground state dye concentration. The similarity of this value to the concentration of triplet formed in the absence of reducing agents and the observation that  $2 \times 10^{-4}$  M ascorbic acid completely quenches the triplet indicate that over the entire pH range, conversion of triplet neutral red to the semireduced radical is essentially quantitative.

Several aspects of the formation of the semireduced radical by ascorbic acid are significant. First, addition of this reducing agent does not decrease the relative yield of neutral red fluorescence, a result which eliminates singlet state neutral red as a direct pathway to the semireduced radical. Second, the total quenching of the triplet transient indicates that conversion of triplet neutral red to the semireduced radical takes place with high efficiency. Even between pH 6 and pH 9, where the triplet decay is so rapid that

it follows the flash lamp decay, the concentration of semireduced neutral red remains at the level observed outside this pH range, where the triplet lifetime is much longer. The ability of the reduction pathway to compete successfully with triplet decay under these extreme circumstances shows clearly that the rate of triplet reduction by 100 fold excess ascorbic acid is much faster than the rate of triplet decay.

In contrast to the highly efficient reduction of triplet neutral red by ascorbic acid from pH 1.3 to pH 11, its reduction by EDTA over this pH range is quite inefficient. For example, over the entire pH interval,  $2 \times 10^{-4}$  M EDTA does not react to a detectable extent with triplet state neutral red. Unfortunately, the low reactivity of EDTA below pH 6 (Bonneau *et al.*, 1974) and the limited solubility of neutral red above pH 7 limit severely the scope of a systematic study of the reduction of triplet state neutral red by EDTA. Transient experiments at pH 6.5 with  $2 \times 10^{-3}$  M EDTA yield small quantities of semireduced dye, equivalent to ca 0.3% of the ground state dye concentration. The low reactivity of EDTA with triplet state neutral red stands in contrast to its high reactivity with the triplet states of thionine, methylene blue (Bonneau *et al.*, 1973; Bonneau *et al.*, 1974) and safranin-O (Baumgartner *et al.*, 1981) both of which EDTA reduces with high efficiency above pH 6. This difference may reflect the fact that the concentration of  ${}^3\text{DH}_2^{+2}$  which exists under these conditions is much lower in the case of neutral


red than in the case of the other dyes. The estimated  $pK_a$ s of  ${}^3DH_2^{+2}$  noted in the previous section support this explanation.

#### Semireduced Radical Spectra

Representative transient spectra of semireduced neutral red secured 1 ms after flash initiation and reported in Figure 3 illustrate the major spectral features of these species. Below pH 6.5, there is a sharp absorption band centered at 390 nm and a broad band centered at 690 nm. Chibisov et al., (1969) reported only the long wavelength band at pH 4. Above pH 8.5 there are two absorption bands at 380 nm and 710 nm. Chibisov et al., (1969), report only a band below 600 nm at pH 11. Neither the wavelengths of the two absorption bands nor the intensity of the shorter wavelength band is sufficiently pH sensitive to be useful for monitoring the acid-base chemistry of semireduced neutral red, but the strong pH dependence of the intensity of the longer wavelength band is ideally suited for this purpose. The normalized absorbance-pH plot at 690 nm reported in Figure 4 displays distinct transitions centered at pH 2.6 and pH 7.8, each of which corresponds to a  $pK_a$ . Chibisov et al., (1969), report a single  $pK_a$  of about 7, but made no measurements below pH 4. Because the rate constant for decay of semireduced neutral red is pH independent (see following section), these transitions can be associated directly with the two  $pK_a$ s of semireduced neutral red. Thus, the acid-base chemistry of semireduced neutral red can be described by Equation 2 in which the



formula of the most highly protonated species is denoted  ${}^3\text{DH}_3^{+2}$ .





This choice of notation is suggested by the similarity of the second  $\text{pK}_a$  of the neutral red radical (7.8) to the  $\text{pK}_a$  of the species  $\text{DH}_2^+$  of thionine (8.2, Fisher, 1964; Bonneau *et al.*, 1968; Faure *et al.*, 1967), oxonine (7.6, Michaelis and Granick, 1941) and safranin-O (9.5, Baumgartner *et al.*, 1981). Molar absorptivities, absorption maxima, and decay constants of the three neutral red radical species along with their  $\text{pK}_a$ s are summarized in Table II. As a check on the correctness of the  $\text{pK}_a$  values, the solid line in Figure 4 is the theoretical line constructed from the  $\text{pK}_a$ s and the limiting absorbances. The close agreement of this theoretical line with the experimental points confirms that these  $\text{pK}_a$  values are consistent with the data.

#### Semireduced Radical Decay

Over the entire pH range, the decay of neutral red formed by ascorbic acid reduction follows second order kinetics with a rate constant of  $1.8 \pm 0.4 \times 10^8 \text{ M}^{-1} \text{ s}^{-1}$ , independent of pH. The two simple processes consistent with these kinetics are the disproportionation of the semireduced radical and the back transfer of an electron from the semireduced radical to oxidized ascorbic acid. The corresponding reactions are defined by Equation 3 and Equation 4, respectively, where AA

and  $AA^+$  represent ascorbic acid and its one electron oxidation product. Although Equations 3 and 4 apply specifically to the pH range 2.6-7.8, similar expressions involving  $DH_3^{+2}$  and  $DH^{\cdot}$  apply at higher and lower pH values.





There are two pieces of evidence which indicate that Equation 3 is not an important pathway for decay of semireduced neutral red. The first is that from pH 1.3 to pH 11 there is no detectable deficit in the concentration of ground state neutral red within 20 ms after subjecting the neutral red-ascorbic acid system to 50 successive flashes, indicating that no ground state neutral red is permanently converted to leuco dye. Since leuco neutral red is reported to be oxidized only slowly near neutral pH on the time scale of potentiometric studies (Clark and Perkins, 1932), the rapid and complete regeneration of ground state neutral red in this pH range in the present study suggests that formation of leuco dye is not significant.

The second and more direct indication that Equation 3 is not the major pathway for the disappearance of semireduced neutral red when a reversible reducing agent is used is the strong dependence of the rate constant for this process on the

identity of the reducing agent. For example, at pH 7.5 with p-phenylenediamine as the reductant, the decay constant of the semireduced radical is  $2.5 \times 10^9 \text{ M}^{-1} \text{ s}^{-1}$ , a value more than 10 fold higher than that observed with ascorbic acid as reductant. Under these conditions the one-electron oxidation product of p-phenylenediamine is a significantly better oxidant than the one-electron oxidation product of ascorbic acid, (Steenken and Neta, 1979). The fact that semireduced neutral red decays ca tenfold more rapidly in the presence of oxidized p-phenylenediamine than in the presence of oxidized ascorbic acid is consistent with the back electron transfer mechanism of Equation 4, but rules out the radical disproportionation mechanism of Equation 3.

Because radical disproportionation is an efficient process for the disappearance of the semireduced radical species of a number of dyes, we considered the more complex mechanism defined by Equations 5 and 6 in which Ox and Re denote the



oxidized and reduced forms of the reducing agent. In this mechanism, a fast radical disproportionation step is followed by a slower step which converts leuco dye to semireduced radical. Because coupling the two steps complicates

analytical solution of the kinetic equations, we used digital simulation to obtain numerical solutions under appropriate conditions. Figure 5 demonstrates the influence of the rate of the coupled reaction between the oxidized form of the reducing agent and the leuco dye on the second order rate plot for disappearance of the semireduced radical. In this simulation,  $k_1$ , the rate constant of Equation 5 is set at  $2 \times 10^8 \text{ M}^{-1} \text{ s}^{-1}$ , while  $k_2$  is varied from 0 to  $2 \times 10^{10} \text{ M}^{-1} \text{ s}^{-1}$ .

As anticipated, the second order kinetic plots are linear if  $k_2$  is either much larger than  $k_1$  or much smaller than  $k_1$ . However, for either condition, the slope of the second order kinetics plot is nearly independent of the value of  $k_2$ . That is, under the conditions which give reasonable second order kinetics plots, the apparent rate constant for disappearance of the semireduced radical is virtually insensitive to the reactivity of the oxidized form of the reducing agent. In fact, over the entire range of  $k_2$ , the value of the apparent second order rate constant varies by only a factor of two. These predictions of the model are clearly inconsistent with the tenfold increase in the rate of disappearance of semireduced dye when ascorbic acid is replaced with p-phenylenediamine, ruling out the disproportionation-regeneration mechanism of Equations 5 and 6. Thus, the only mechanism consistent with all of the kinetic data is the back electron transfer process of Equation 4.

## ACKNOWLEDGEMENT

This work was supported in part by the Office Of Naval Research. The helpful advice of Dr. Ekehardt Vogelmann is gratefully acknowledged.

## REFERENCES

- Bartels, P. (1956) Z. Phys. Chem. N.F. 9, 74-94.
- Baumgartner, C.E., H.H. Richtol and D.A. Aikens (1981) Photochem. Photobiol. 34, 17-22.
- Bishop, E. (1972) Indicators, pp. 490-493. Pergamon Press Oxford.
- Bonneau, R. And J. Joussot-Dubien (1973) Photochem. Photobiol. 17, 313-319.
- Bonneau, R., P. Fornier de Violet and J. Joussot-Dubien (1974) Photochem. Photobiol. 19, 129-132.
- Bonneau, R. and J. Pereyre (1975) Photochem. Photobiol. 21, 173-177.
- Bonneau, R., J. Faure and J. Joussot-Dubien (1981). Ber. Bunsenges. Phys. Chem. 72, 263-266.
- Chibisov, A.K., V.B. Skvortsov, A.V. Karyakin, and L.N. Rygalov (1969) Khim. Vys. Energ. 3, 210-216.
- Clark, W.M. and M.E. Perkins (1932) J. Am. Chem. Soc. 54, 1228-1247.
- Faure, J., R. Bonneau, and J. Joussot-Dubien (1965) J. Chim. Phys. 65, 369-370.
- Faure, J., R. Bonneau and J. Joussot-Dubien (1967) Photochem. Photobiol. 6, 331-339.
- Fischer, H. (1964) Z. Phys. Chem. N.F. 43, 177-194.
- Knowles, A. (1971) Photochem. Photobiol. 13, 473-478.
- Michaelis, L. and S. Granick (1941) J. Am. Chem. Soc. 63, 1636-1643.
- Ohno, T. and N.N. Lichtin (1980) J. Am. Chem. Soc. 102, 4636-4643.
- Schulman, S.G. (1976) in Modern Fluorescence Spectroscopy (Edited by E.L. Wehry), pp. 239-275. Plenum Press, New York.
- Steenken, S. And P. Neta (1979) J. Phys. Chem. 83, 1134-1137.

Strong, R.L. And J. Perano (1967) J. Am. Chem. Soc. 89,  
2535-2538.

TABLE I  
Properties of Neutral Red Triplet Species

<u>Species</u>	<u>pK<sub>a</sub></u>	<u>Absorption Maxima (nm)</u>	<u>Molar Absorptivity (M<sup>-1</sup>cm<sup>-1</sup>)</u>	<u>Decay Constant (s<sup>-1</sup>)</u>
<sup>3</sup> DH <sub>2</sub> <sup>+2</sup>		390 680	8,000 12,000	1.6 ± 0.3 × 10 <sup>4</sup>
<sup>3</sup> DH <sup>+</sup>	4.8			
	9.5			
<sup>3</sup> D		380 580	5,800 7,300	1.6 ± 0.3 × 10 <sup>4</sup>




TABLE II  
Properties of Semireduced Neutral Red Species


<u>Species</u>	<u>pK<sub>a</sub></u>	<u>Absorption Maxima (nm)</u>	<u>Molar Absorptivity (M<sup>-1</sup>cm<sup>-1</sup>)</u>	<u>Decay Constant (M<sup>-1</sup>s<sup>-1</sup>)</u>
DH <sub>3</sub> <sup>+2</sup>	2.6	390	8200	1.8 ± 0.4 × 10 <sup>8</sup>
		690	1000	
DH <sub>2</sub> <sup>+</sup>	7.8	390	8200	1.8 ± 0.4 × 10 <sup>8</sup>
		690	5800	
DH <sup>•</sup>		380	7000	1.8 ± 0.4 × 10 <sup>8</sup>
		710	8100	


## FIGURE CAPTIONS


- Figure 1    Transient absorption spectra of triplet species  
              open circles - pH 2.8  
              closed circles - pH 9.8
- Figure 2    Normalized absorbance - pH plot for triplet species  
              open circle - acidic triplet at 680 nm  
              closed circles - basic triplet at 580 nm
- Figure 3    Transient absorption spectra of semireduced radicals  
              open circles - pH 2.8  
              closed circles - pH 9.8
- Figure 4    Normalized absorbance-pH plot for semireduced radicals at 690 nm
- Figure 5    Digitally simulated second order plots for disappearance of semireduced radical by mechanism of Eqns 5 and 6 with  $k_1 = 2 \times 10 \text{ M}^{-1} \text{ s}^{-1}$ .


plot	$k_2 \text{ M}^{-1} \text{ s}^{-1}$
a	0
b	$2 \times 10^7$
c	$6 \times 10^7$
d	$2 \times 10^8$
e	$6 \times 10^8$
f	$2 \times 10^{10}$











TECHNICAL REPORT DISTRIBUTION LIST, GEN

	<u>No. Copies</u>		<u>No. Copies</u>
Office of Naval Research Attn: Code 413 800 North Quincy Street Arlington, Virginia 22217	2	Naval Ocean Systems Center Attn: Mr. Joe McCartney San Diego, California 92152	1
ONR Pasadena Detachment Attn: Dr. R. J. Marcus 1030 East Green Street Pasadena, California 91106	1	Naval Weapons Center Attn: Dr. A. B. Amster, Chemistry Division China Lake, California 93555	
Commander, Naval Air Systems Command Attn: Code 310C (H. Rosenwasser) Department of the Navy Washington, D.C. 20360	1	Naval Civil Engineering Laboratory Attn: Dr. R. W. Drisko Port Hueneme, California 93401	1
Defense Technical Information Center Building 5, Cameron Station Alexandria, Virginia 22314	12	Dean William Tolles Naval Postgraduate School Monterey, California 93940	1
Dr. Fred Saalfeld Chemistry Division, Code 6100 Naval Research Laboratory Washington, D.C. 20375	1	Scientific Advisor Commandant of the Marine Corps (Code RD-1) Washington, D.C. 20380	1
U.S. Army Research Office Attn: CRD-AA-IP P. O. Box 12211 Research Triangle Park, N.C. 27709	1	Naval Ship Research and Development Center Attn: Dr. G. Bosmajian, Applied Chemistry Division Annapolis, Maryland 21401	1
Mr. Vincent Schaper DTNSRDC Code 2803 Annapolis, Maryland 21402	1	Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 19112	1
Naval Ocean Systems Center Attn: Dr. S. Yamamoto Marine Sciences Division San Diego, California 91232	1	Mr. A. M. Anzalone Administrative Librarian PLASTEC/ARRADCOM Bldg 3401 Dover, New Jersey 07801	1



TECHNICAL REPORT DISTRIBUTION LIST, 359

	<u>No. Copies</u>		<u>No. Copies</u>
Dr. Paul Delahay Department of Chemistry New York University New York, New York 10003	1	Dr. P. J. Hendra Department of Chemistry University of Southampton Southampton SO0 5NH United Kingdom	1
Dr. E. Yeager Department of Chemistry Case Western Reserve University Cleveland, Ohio 41106	1	Dr. Sam Perone Chemistry & Materials Science Department Laurence Livermore National Lab. Livermore, California 94550	1
Dr. D. N. Bennion Department of Chemical Engineering Brigham Young University Provo, Utah 84602	1	Dr. Royce W. Murray Department of Chemistry University of North Carolina Chapel Hill, North Carolina 27514	1
Dr. R. A. Marcus Department of Chemistry California Institute of Technology Pasadena, California 91125	1	Naval Ocean Systems Center Attn: Technical Library San Diego, California 92152	1
Dr. J. J. Auburn Bell Laboratories Murray Hill, New Jersey 07974	1	Dr. C. E. Mueller The Electrochemistry Branch Materials Division, Research and Technology Department Naval Surface Weapons Center White Oak Laboratory Silver Spring, Maryland 20910	1
Dr. Adam Heller Bell Laboratories Murray Hill, New Jersey 07974	1	Dr. G. Goodman Johnson Controls 5757 North Green Bay Avenue Milwaukee, Wisconsin 53201	1
Dr. T. Katan Lockheed Missiles and Space Co., Inc. P. O. Box 504 Sunnyvale, California 94088	1	Dr. J. Boechler Electrochimica Corporation Attn: Technical Library 2485 Charleston Road Mountain View, California 94040	1
Dr. Joseph Singer, Code 302-1 NASA-Lewis 21000 Brookpark Road Cleveland, Ohio 44135	1	Dr. P. P. Schmidt Department of Chemistry Oakland University Rochester, Michigan 48063	1
Dr. B. Brummer EIC Incorporated 55 Chapel Street Newton, Massachusetts 02158	1		
Library P. R. Mallory and Company, Inc. Northwest Industrial Park Burlington, Massachusetts 01803	1		

TECHNICAL REPORT DISTRIBUTION LIST, 359

	<u>No. Copies</u>		<u>No. Copies</u>
Dr. H. Richtol Chemistry Department Rensselaer Polytechnic Institute Troy, New York 12181	1	Dr. R. P. Van Duyne Department of Chemistry Northwestern University Evanston, Illinois 60201	1
Dr. A. B. Ellis Chemistry Department University of Wisconsin Madison, Wisconsin 53706	1	Dr. B. Stanley Pons Department of Chemistry University of Alberta Edmonton, Alberta CANADA T6G 2G2	1
Dr. M. Wrighton Chemistry Department Massachusetts Institute of Technology Cambridge, Massachusetts 02139	1	Dr. Michael J. Weaver Department of Chemistry Michigan State University East Lansing, Michigan 48824	1
Larry E. Plew Naval Weapons Support Center Code 30736, Building 2906 Crane, Indiana 47522	1	Dr. R. David Rauh EIC Corporation 55 Chapel Street Newton, Massachusetts 02158	1
S. Ruby DOE (STOR) 600 E Street Providence, Rhode Island 02192	1	Dr. J. David Margerum Research Laboratories Division Hughes Aircraft Company 3011 Malibu Canyon Road Malibu, California 90265	1
Dr. Aaron Wold Brown University Department of Chemistry Providence, Rhode Island 02192	1	Dr. Martin Fleischmann Department of Chemistry University of Southampton Southampton 509 5NH England	1
Dr. R. C. Chudacek McGraw-Edison Company Edison Battery Division Post Office Box 28 Bloomfield, New Jersey 07003	1	Dr. Janet Osteryoung Department of Chemistry State University of New York at Buffalo Buffalo, New York 14214	1
Dr. A. J. Bard University of Texas Department of Chemistry Austin, Texas 78712	1	Dr. R. A. Osteryoung Department of Chemistry State University of New York at Buffalo Buffalo, New York 14214	1
Dr. M. M. Nicholson Electronics Research Center Rockwell International 3370 Miraloma Avenue Anaheim, California	1		

TECHNICAL REPORT DISTRIBUTION LIST, 359

	<u>No. Copies</u>		<u>No. Copies</u>
Dr. Donald W. Ernst Naval Surface Weapons Center Code R-33 White Oak Laboratory Silver Spring, Maryland 20910	1	Mr. James R. Moden Naval Underwater Systems Center Code 3632 Newport, Rhode Island 02840	1
Dr. R. Nowak Naval Research Laboratory Code 6130 Washington, D.C. 20375	1	Dr. Bernard Spielvogel U. S. Army Research Office P. O. Box 12211 Research Triangle Park, NC 27709	1
Dr. John F. Houlihan Shenango Valley Campus Pennsylvania State University Sharon, Pennsylvania 16146	1	Dr. Denton Elliott Air Force Office of Scientific Research Bolling AFB Washington, D.C. 20332	1
Dr. D. F. Shriver Department of Chemistry Northwestern University Evanston, Illinois 60201	1	Dr. David Aikens Chemistry Department Rensselaer Polytechnic Institute Troy, New York 12181	1
Dr. D. H. Whitmore Department of Materials Science Northwestern University Evanston, Illinois 60201	1	Dr. A. P. B. Lever Chemistry Department York University Downsview, Ontario M3J1P3 Canada	1
Dr. Alan Bewick Department of Chemistry The University Southampton, SO9 5NH England	1	Dr. Stanislaw Szpak Naval Ocean Systems Center Code 6343 San Diego, California 95152	1
Dr. A. Himy NAVSEA-5433 NC #4 2541 Jefferson Davis Highway Arlington, Virginia 20362	1	Dr. Gregory Farrington Department of Materials Science and Engineering University of Pennsylvania Philadelphia, Pennsylvania 19104	1
Dr. John Kincaid Department of the Navy Strategic Systems Project Office Room 901 Washington, D.C. 20376	1	Dr. Bruce Dunn Department of Engineering & Applied Science University of California Los Angeles, California 90024	1

TECHNICAL REPORT DISTRIBUTION LIST, 359

	<u>No. Copies</u>		<u>No. Copies</u>
M. L. Robertson Manager, Electrochemical and Power Sonics Division Naval Weapons Support Center Crane, Indiana 47522	1	Dr. T. Marks Department of Chemistry Northwestern University Evanston, Illinois 60201	1
Dr. Elton Cairns Energy & Environment Division Lawrence Berkeley Laboratory University of California Berkeley, California 94720	1	Dr. D. Cipris Allied Corporation P. O. Box 3000R Morristown, New Jersey 07960	1
Dr. Micha Tomkiewicz Department of Physics Brooklyn College Brooklyn, New York 11210	1	Dr. M. Philpot IBM Corporation 5600 Cottle Road San Jose, California 95193	1
Dr. Lesser Blum Department of Physics University of Puerto Rico Rio Piedras, Puerto Rico 00931	1	Dr. Donald Sandstrom Washington State University Department of Physics Pullman, Washington 99164	1
Dr. Joseph Gordon, II IBM Corporation K33/281 5600 Cottle Road San Jose, California 95193	1	Dr. Carl Kannewurf Northwestern University Department of Electrical Engineering and Computer Science Evanston, Illinois 60201	1
Dr. Robert Somoano Jet Propulsion Laboratory California Institute of Technology Pasadena, California 91103	1	Dr. Edward Fletcher University of Minnesota Department of Mechanical Engineering Minneapolis, Minnesota 55455	1
Dr. Johann A. Joebstl USA Mobility Equipment R&D Command DRDME-EC Fort Belvoir, Virginia 22060	1	Dr. John Fontanella U.S. Naval Academy Department of Physics Annapolis, Maryland 21402	1
Dr. Judith H. Ambrus NASA Headquarters M.S. RTS-6 Washington, D.C. 20546	1	Dr. Martha Greenblatt Rutgers University Department of Chemistry New Brunswick, New Jersey 08903	1
Dr. Albert R. Landgrebe U.S. Department of Energy M.S. 6B025 Forrestal Building Washington, D.C. 20595	1	Dr. John Wassib Kings Mountain Specialties P. O. Box 1173 Kings Mountain, North Carolina 28086	1

TECHNICAL REPORT DISTRIBUTION LIST, 359

	<u>No. Copies</u>	<u>No. Copies</u>
Dr. J. J. Brophy University of Utah Department of Physics Salt Lake City, Utah 84112	1	
Dr. Walter Roth Department of Physics State University of New York Albany, New York 12222	1	
Dr. Thomas Davis National Bureau of Standards Polymer Science and Standards Division Washington, D.C. 20234	1	
Dr. Charles Martin Department of Chemistry Texas A&M University	1	
Dr. Anthony Sammells Institute of Gas Technology 3424 South State Street Chicago, Illinois 60616	1	
Dr. H. Tachikawa Department of Chemistry Jackson State University Jackson, Mississippi 39217	1	
Dr. W. M. Risen Department of Chemistry Brown University Providence, Rhode Island	1	

**END**

**FILMED**

**9-83**

**DTIC**